

Let's talk about Software

GROSTAT VI, Menton, February 20, 2003

Raymond Hemmecke

UC Davis

raymond@hemmecke.de

One Problem

Sampling/moves on 3-way tables – Markov bases

0	1	0	1	1	0	1
1	0	1	0	1	0	1
0	1	1	0	0	1	1
1	1	0	0	0	1	1
0	0	0	0	1	1	0
0	0	1	1	1	1	0

Markov basis \subseteq Gröbner basis \subseteq Graver basis

One Answer

4_{ti}2

by R. & R. Hemmecke

downloadable from www.4ti2.de

4_{ti}2


- (truncated) **Hilbert bases**
 - 6×6 magic squares ($\sim 500,000$ elements, ~ 10 days)
- (truncated, symmetric) **Graver bases**
 - hppi's of order 20 ($\sim 1,250,000$ elements, ~ 1 month)
- (symmetric) **Gröbner bases** for graded toric ideals
 - $3 \times 4 \times 5$ tables ($\sim 77,000$ elements, ~ 4 days)
 - binary K_5 model ($\sim 13,000$ elements, ~ 30 hours)

Challenges for $4_{ti}2$

- redo $4 \times 4 \times 4$ tables (Sawae et al.)
- Sullivant: verify that a given set of 145,512 binomials generate the toric ideal for $4 \times 4 \times 4$ tables
- Hosten: binary K_6 model and its submodels
- Sturmfels: binary octahedron model
- $3 \times 3 \times 3 \times 3$ tables


Let's do some magic with $4_{ti}2$

Hilbert bases and magic squares



Lemma: The set of non-negative real magic squares forms a cone.

What is a Hilbert basis?



$(0, 3, 1)$ $(2, 2, 1)$ $(4, 1, 1)$ $(6, 0, 1)$

Solving the problem with $4_{ti}2$

Consider the pointed rational cone of magic squares:

$$\ker(A) \cap \mathbb{R}_+^n = \{z : Az = 0, z \in \mathbb{R}_+^n\}.$$

Every row, column and diagonal sum equals the sum of the first row.

Input file: 3x3magic

9 7

```
1 1 1 -1 -1 -1 0 0 0
1 1 1 0 0 0 -1 -1 -1
0 1 1 -1 0 0 -1 0 0
1 0 1 0 -1 0 0 -1 0
1 1 0 0 0 -1 0 0 -1
0 1 1 0 -1 0 0 0 -1
1 1 0 0 -1 0 -1 0 0
```

Call $4_{ti}2$

```
./hilbert 3x3magic
```

Output file: 3x3magic.hil

```
9 5  
1 2 0 0 1 2 2 0 1  
0 2 1 2 1 0 1 0 2  
1 1 1 1 1 1 1 1 1  
2 0 1 0 1 2 1 2 0  
1 0 2 2 1 0 0 2 1
```

Hilbert basis: 3×3 magic squares

1	0	2
2	1	0
0	2	1

2	0	1
0	1	2
1	2	0

0	2	1
2	1	0
1	0	2

1	2	0
0	1	2
2	0	1

1	1	1
1	1	1
1	1	1

Generating Function of 3×3 Magic Squares

Generating function of 3×3 magic squares

$$g_3(t) = \frac{(t^3 + 1)^2}{(1 - t^3)^3} = \sum \left(\begin{array}{c} \#3 \times 3 \\ \text{magic squares} \\ \text{with sum } s \end{array} \right) t^s$$

The number of 3×3 magic squares of sum s is

$$M_3(s) = \begin{cases} \frac{2}{9}s^2 + \frac{2}{3}s + 1 & \text{if } 3|s, \\ 0 & \text{otherwise,} \end{cases}$$

This final series computation was done by **CoCoA**.

New Formulas for Magic and Semi-Magic Cubes

Theorem [Ahmed, De Loera, H.] The number of $3 \times 3 \times 3$ “semi-magic” cubes is

$$\begin{cases} \frac{9}{2240} s^8 + \frac{27}{560} s^7 + \frac{87}{320} s^6 + \frac{297}{320} s^5 + \frac{1341}{640} s^4 + \frac{513}{160} s^3 + \frac{3653}{1120} s^2 + \frac{627}{280} s + 1 & \text{if } 2|s, \\ \frac{9}{2240} s^8 + \frac{27}{560} s^7 + \frac{87}{320} s^6 + \frac{297}{320} s^5 + \frac{1341}{640} s^4 + \frac{513}{160} s^3 + \frac{3653}{1120} s^2 + \frac{4071}{2240} s + \frac{47}{128} & \text{otherwise.} \end{cases}$$

Theorem[Ahmed, De Loera, H.] The number of $3 \times 3 \times 3$ magic cubes is

$$MC_3(s) = \begin{cases} \frac{11}{324} s^4 + \frac{11}{54} s^3 + \frac{25}{36} s^2 + \frac{7}{6} s + 1 & \text{if } 3|s, \\ 0 & \text{otherwise.} \end{cases}$$

Computations were done by **4ti2** and **CoCoA**.

What is a Graver basis?

What is a Graver basis?

$$G(A) = \bigcup_j H(\ker(A) \cap \mathbb{O}_j)$$

What is a Graver basis?

$$G(A) = \bigcup_j H(\ker(A) \cap \mathbb{O}_j)$$

Fact: The Graver basis of A contains the **Markov basis** of A .

3×3 tables

9 6

1	1	1	0	0	0	0	0	0
0	0	0	1	1	1	0	0	0
0	0	0	0	0	0	1	1	1
1	0	0	1	0	0	1	0	0
0	1	0	0	1	0	0	1	0
0	0	1	0	0	1	0	0	1

Call $4t_2$

`./graver 3x3`

or

`./groebner 3x3`

Output file: 3x3.gra

9 15

```
-1  1  0  1 -1  0  0  0  0
 0  0  0  1  0 -1 -1  0  1
-1  0  1  1  0 -1  0  0  0
 1 -1  0  0  0  0 -1  1  0
 0  0  0  1 -1  0 -1  1  0
-1  1  0  1  0 -1  0 -1  1
 0 -1  1  1  0 -1 -1  1  0
 1 -1  0  0  1 -1 -1  0  1
-1  0  1  0  0  0  1  0 -1
```

...

Let us exploit symmetry

Define symmetry group: 3x3.sym

```
9 4  
2 3 1 5 6 4 8 9 7  
2 1 3 5 4 6 8 7 9  
4 5 6 7 8 9 1 2 3  
4 5 6 1 2 3 7 8 9
```

Call **4_{ti}2** with symmetry

```
./graver symmetry 3x3
```

or just

```
./graver sym 3x3
```

which creates a file **3x3.sgra**

Call $4_{ti}2$ with symmetry

```
./graver symmetry 3x3
```

or just

```
./graver sym 3x3
```

which creates a file 3x3.sgra and then **type**

```
./output 3way 3 3 1 3x3.sgra
```

which creates a file **3x3.sgra.3way**

Output file: 3x3.sgra.3way

3 3 1

=====

0 0 0
1 0 -1
-1 0 1

=====

0 1 -1
1 -1 0
-1 0 1

=====

Two representatives up to symmetry

$$\begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ -1 & 0 & 1 \end{pmatrix}$$

$3 \times 3 \times 4$ tables

	Gröbner	symmGröbner	Graver	symmGraver
Elements	626	9	19,722	37
Time	47	447	1978	437

Another Problem

Is there a table?

?	?	?	?	220
?	?	?	?	215
?	?	?	?	93
?	?	?	?	64
108	286	71	127	

There is an integer solution to $Ax = b, x \geq 0$ if and only if there is a minimal solution to $(A| -b)y = 0, y \geq 0$ with last component 1.

Is there a table?

?	?	?	?	220
?	?	?	?	215
?	?	?	?	93
?	?	?	?	64
108	286	71	127	

There is an integer solution to $Ax = b, x \geq 0$ if and only if there is a minimal solution to $(A| -b)y = 0, y \geq 0$ with last component 1.

In this way, you may find that Hosten's $2 \times 3 \times 3$ table has only 441 solutions.

How many tables are there?

68	119	26	7	220
20	84	17	94	215
15	54	14	10	93
5	29	14	16	64
108	286	71	127	

How many tables are there?

68	119	26	7	220
20	84	17	94	215
15	54	14	10	93
5	29	14	16	64
108	286	71	127	

There are 1,225,914,276,768,514 such tables.

How many tables are there???

?	?	?	?	?	338106
?	?	?	?	?	574203
?	?	?	?	?	678876
?	?	?	?	?	1213008

2
0
2
0
2

1
4
2
7
4
6

4
1
0
7
5
5

1
0
0
7
7
7
3

1
2
2
2
7
1
7

Another Answer

LattE - Counting lattice points

- software package developed at UC Davis (De Loera, H., Tauzer, Yoshida)
- computes number of lattice points in **any** rational polytope

$$Ax = a, Bx \geq b$$

- includes several algorithms to count lattice points

www.math.ucdavis.edu/~latte

One way to count

One way to count

Define generating function

$$g_P(z) = \sum_{\alpha \in P} z^\alpha$$

One way to count

Define generating function

$$g_P(z) = \sum_{\alpha \in P} z^\alpha$$

Then the number of lattice points is

$$g_P(\mathbf{1}) = \sum_{\alpha \in P} 1$$

Life is not always easy

A closer look onto this approach

Problem: $g_P(z)$ written as a polynomial is far too big.

A closer look onto this approach

Problem: $g_P(z)$ written as a polynomial is far too big.

Solution: Write $g_P(z)$ as a sum of rational functions $p(z)/q(z)$.

A closer look onto this approach

Problem: $g_P(z)$ written as a polynomial is far too big.

Solution: Write $g_P(z)$ as a sum of rational functions $p(z)/q(z)$.

Barvinok: There is always a short (polynomial size) representation of $g_P(z)$.

Problems solved

- recomputation of counts from literature
- recomputation of counting formulas for $3 \times 3 \times 3$ magic and semi-magic cubes
- Aardal's Frobenius problems
- $3 \times 3 \times 4$ tables for given marginals
- some new counting formulas (24-cell, . . .)

Challenges for LattE

- new counting formulas for $4 \times 4 \times 4$ magic and semi-magic cubes
- new counting formula for 5×5 magic squares
- your problems?!

4_{ti}2

www.4ti2.de

LattE

www.math.ucdavis.edu/~latte